

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of Men, and that it is frequently made use of for the taking of Films from the Eyes of Horses and other Beasts. I hear also that they sell good quantities of it to the Druggists of London, but for what use I know not. This is all I could learn concerning Calamine. If I have omitted any thing wherein you are desirous of surther Information; or if in any other Concern of this Nature I can be serviceable, you may freely command,

Wrington, Oct.25.84.

Tours, &c.

VIII. An Arithmetical Paradox, concerning the Chances of Lotteries, by the Honourable Francis Roberts, Esq. Fellow of the R. S.

S some Truths (like the Axiomes of Geometry and Metaphysicks) are self-evident at the first View, to there are others no less certain in their Foundation, that have a very different Aspect, and without a strict and careful Examination rather seem repugnant.

We may find Instances of this kind in most Sci-

ences.

In Geometry, That a Body of an infinite Length may yet have but a finite Magnitude.

In Geography, That if Antwerp be due East to London, for that reason London cannot be West to Ant-

werp.

In Astronomy, That at the Barbadoes (and other places between the Line and Tropick) the Sun, part of the Year, comes twice in a Morning to some Points of the Compass.

In Hydrostaticks, That a hollow Cone (standing upon its Basis) being fill'd with Water, the Water shall press the bottom with three times the Weight, as if the same Water was frozen to Ice; and Figures might be contrived to make it press a hundred times as much.

These Speculations, as they are generally pleasant, so they may also be of good use to warn us of the Mistakes we are liable to, by careless and superficial reason-

ing.

I shall add one Instance in Arithmetick, which perhaps may seem as great a Paradox as any of the former.

There are two Lotteries, at either of which a Gamefler paying a Shilling for a Lot or Throw; the First Lottery upon a just Computation of the Odds has 3 to 1 of the Gamester, the Second Lottery but 2 to 1; nevertheless the Gamester has the very same disadvantage (and no more) in playing at the First Lottery as the Second.

It looks very like a Contradiction, that the Disadvantage should be no greater in playing against 3 to 1 than 2 to 1, but it may thus be resolv'd.

Let the
$$\left\{\begin{array}{c} 1/h \\ 2d \end{array}\right\}$$
 Lottery con- $\left\{\begin{array}{c} 3 \\ 4 \\ \end{array}\right\}$ Blanks $\left\{\begin{array}{c} 3 \\ 2 \\ \end{array}\right\}$ Prizes $\left\{\begin{array}{c} 16 \text{ pence} \\ 2 \text{ fhill.} \end{array}\right\}$ a piece.

In the first Lottery the Gamester hazards a Shilling to win a Groat, and the Chances being equal, it is evident there is 3 to one against him.

In the Second Lottery the Gamester ventures a Shilling against a Shilling, and the Lots being 4 to 2, his

Disadvantage is 2 to 1.

And a Lot at either of them being truly worth just 8 Pence, (viz. the 6th part of 3 times 16 Pence, or twice 2 Shillings) the Disadvantage must be the very

same in both Cases, that is, the Gamester pays a Shil-

ling for a Lot that is worth but 8 Pence.

The Method of finding this Answer being somewhat out of the common Road, I shall here add it, and thereby infinite Solutions of the same kind may be discovered.

Ist Lottery.

2d Lottery.

Let a= the number of Blanks. m= the number of Blanks b= the number of Prizes. n= the Value of a Prize. s= the value of a Prize.

1 = to what you pay for a Lot, viz. a Shilling.

So the Lottery has it's Chances for 1, and the Gamefter his for r-1. Now the true Odds confisting of the compounded Proportion of the Chances and the Values, viz. $\frac{a}{b}$ and $\frac{1}{r-1}$, the Share of the Lottery will be a, and that of the Gamester rb-b. Therefore as the present case stands, the first Lottery must be a=3 rb-3b, and by the like reasoning the second Lottery will be m=2 s n-2 n. Now the Value of a Lot being the Sum of the Prizes divided by the number of Lots, (which must be equal in both Lotteries) it yields

$$\frac{r\,b}{a+b} = \frac{s\,n}{m+n}$$

So to proceed.

$$\begin{bmatrix} a \\ b \\ r \\ s \end{bmatrix} = \begin{vmatrix} 1 \\ 2 \\ m = 2 \\ s \\ n - 2 \\ n \\ 4 \\ (*) \\ (*) \\ 6 \\ (*) \end{vmatrix}$$

```
(680)
 q = \frac{?}{|a|} |7| \operatorname{Let} \frac{rb}{a+b} = q
 7 * \overline{a+b} \begin{vmatrix} 8 & rb = qa + qb \\ 8 \times \overline{3} \end{vmatrix} 9 \begin{vmatrix} 3rb = 3qa + 3qb \end{vmatrix}
   |11|39a+39b=a+3b
   9, 10
              12 If a = 0 to avoid negative Numbers.
   Scope
              |13|3b = 3qb
 II, I2
 13 ÷ 3b
               14 q = 1
              15 q > 1 makes a < 0 q < 1 makes a > 0
 12, 14
              16 If b =0
 Scope
              |17|3qa=a
 11, 16
               18 | q = \frac{1}{4}
 17 ÷ 3a
 16, 18
               19 | q < \frac{1}{2}  makes b < 0 q > \frac{1}{2} makes b > 0
       7,
20*m-n
               2I \mid Sn = qm + qn
               22 | 25n = 2qm + 2qn
 2I * 2
 2+2n
               23 | 25n = m + 2n
               24 | 2qm + 2qn = m + 2n
 22, 23
               25 | \text{If } m = 0
 Scope
              |26|29n = 2n
 24, 25
              |27|q = 1
 26 ÷ 2 n
                    q > 1 makes m < 0 g < 1 makes m > 0
 25, 27
              |29| \text{If } n = 0
 Scope
               30 | 2qm = m
 24, 29
 30 \div 2 \ m \ | \ 31 \ | \ q = \frac{1}{2}
               32 \left| q \right| < \frac{1}{2} \text{ makes } n < 0 \quad q > \frac{1}{2} n > 0
 29, 31
               33 that a b m n \text{ may be} > 0, q \text{ must be} > \frac{1}{2} < r
```

33, 4 (*)

7, 34.

35
$$\frac{r}{a+b} = \frac{2}{3}$$

36 $\frac{rb}{a+b} = \frac{2}{3}$

37 $\frac{rb}{a+b} = \frac{2}{3}$

38 * 39 $\frac{rb}{a+b} = \frac{2}{3}$

38 * 39 $\frac{rb}{a+b} = \frac{2}{3}$

39 * 2 40 6sn = 2m + 2n

6sn = 3m + 6n

40, 41 41 4m + 4n = 3m + 6n

42 - 43 $m = 2n$

1 ÷ 37 44 1 = 3r - 3

44 + 3 2 = 2s - 2

5 (*)

48 Let $A = 3$

37, 48 49 $B = 3$

49 $B = 3$

50 $R = \frac{4}{3}$, id eft, 16 Pence.

Let $M = 4$

43, 51 52 $N = 2$

5 S = 2 2 Shillings.